## 1. REFERENCES


Abirami G. (2014), A New RP-HPLC method for simultaneous estimation of Thiocolchicoside and Ketoprofen in tablet dosage form, World J Pharm & Pharm Sci, 3(2), 2564-75.

Albert R. (1997), Validation of analytical procedure, Ana. Chem. 789.

Altun M. (2002), Turkish Journal of Chemistry, 26, 521-8.

American Heart Association (2009), Heart disease and stroke statistics-2009 Update, A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee Circulation, 119, e21-e181.

Amin M. and Seth K. (2012), Development and Validation of stability indicating assay method for Lornoxicam & Tramadol in Tablet Dosage Form by RP-HPLC, Pharma Science Monitor, Int J Pharm Sci, 3(2), 11-29.

Basaviah K. (2009), Spectrophotometric determination of Lisinopril in pharmaceuticals using Ninhydrin-a modified approach, J of food and drug analysis, 17(2), 93-9.

Bhusari K. and Dhaneshwar R. (2012), Development of a validated Stability-indicating HPLC assay method for Dexketoprofen Trometamol, Int J Pharm Pharm Sci, 4(1), 321-326.

Birajdar S. and Suresh B. (2009), Intentional Journal of Pharmaceutical Research and Development, .2009, 1(1), 23-31.

Boopathy.D. and Perumal.P. (2010), Analytical method development and validation of Losartan Potassium and Atenolol in combined dosage form by RP-HPLC. Int. J. PharmTech Res., 2(01), 471-4.

British Pharmacopoeia, (2009), Medicines and Healthcare Products Regulatory Agency, 512-5.

Chouhan.K., Jain.S. and Subramaniam A. (2011), Validated RP-HPLC method for simultaneous determination of Atorvastatin and Ramipril and its application in drug formulation. Schr. Res. Lbry. Der Pharmacia Lettre, 3(2), 194-202.

Damodiya V., Singh R., Jat K. and Chokshi R. (2012), Method Development and validation of Terbinafine HCl in cream by RP-HPLC, Inventi Impact Pharm Ana & Qual Assurance, 307.

Devi R. and Ramakrishna S. (2010), New spectrophotometric methods for simultaneous estimation of Amlodipine besylate and Atorvastatin calcium in tablet dosage form. Int.J Pharmacy and Pharm Sciences, 2(4), 215.

Dhabale N., Burande B., Hosmani H. and Rakesh U. (2009), Development and statistical validation of UV spectrophotometric method for estimation of Amlodipine besylate in tablet dosage form. Arch pharm sci and res.,1(2),158-61.

Dhaneshwar R. and Bhusari K. (2011), Validated HPTLC method for simultaneous estimation of Thiocolchicoside and Aceclofenac in bulk drug and Formulation, Int J Pharm Bio Sci, 2(2), 482-90.

Dinesh kumar jain et.al International Journal of Pharm Biomed sci 2011,2(2),55-60.

Divyakanth P., Sharad K., Yogesh Y., Seth K., Gajanan D. and Ghelani T. (2011), RP-HPLC methods for development and validation of Lisinopril tablets. An Int.J of pharmaceutical sciences, 1000-8.

Fegade D., Bhole P. and Patil, R. (2009), International Journal of Pharm Tech Res., 1, 184-90.

Franeta T. and Agbaba D. (2002), Il Farmaco, 57(9), 709-13.

Fuad R. (2010), Saudi Pharmaceutical Journal, 2010, 18(4), 103-6.

Garry C. (2000), Analytical Chemistry, University of Washington, 2, 469-75.

Gopal G., Shailendra S. and Swarnlata S. (2008), Development and validation simultaneous estimation of Amlodipine besylate and Enalapril maleate in combined dosage form; Trends in Applied Science Research, 3(3), 278-84.

Gowri N., Vaidhyalingam V. and Shantha A. (2002), Simultaneous estimation of Amlodipine besylate and Benazepril from tablet by RP-HPLC. Indian Drugs, 39(10), 532-35.

Hadad GM, Emara S, Waleed MM, Talanta,.2009, 79(10); 1360-1367.

Hyun L. (2008), Journal of Chromatography, 863(1), 158-162.

ICH-Topic Q2A (2002), Validation of Analytical Procedures, International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use, Geneva.

ICH-Topic Q2B (2002), Validation of Analytical Procedures: Methodology, International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use, Geneva.

ICH-Topic Q6B (2002), Specifications Biotechnological substances, International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use, Geneva.

Jamakhandi M., Chandrasekhar J., Satish K., Santosh K. and Sanjay K. (2010), New fluorimetric method of determination for Lisinopril dosage forms. Int.J of pharma sciences and drug research, 2(3), 182-87.

Kadam A. (2013), Development and Validation of Reversed- Phase High Performance Liquid Chromatographic Method for Estimation of Dexketoprofen Trometamol in Bulk and Tablet Dosage Form, American Journal of Phytomedicine and Clinical Therapeutics, 1(4), 395-400.

Karunakaran K, Navaneethan G. and Pitchaimuthu E. K, (2014), A Validated Stability-Indicating RP-HPLC method for Paracetamol and Lornoxicam : Application to Pharmaceutical Dosage Forms, Chem. Ind. Chem. Eng. Q, 20 (1), 109-114.

Kousy N. (1999), Journal of Pharmaceutical and Biomedical Analysis, 20(6); 185-194.

Kovala D. (2006), Recent advances on non-steroidal anti-inflammatory drugs, NSAIDs: organotin complexes of NSAIDs, Journal of Organometallic Chemistry, 691 (8), 1767–74.

Kumar S. and Prasad R. (2010), Development and validation of reversed phase HPLC method for simultaneous estimation of telmisartan and amlodipine in tablet dosage form. Int. J. Pharm. and Ph'cal. Sci., 2(03), 128-31.

Kumar S., Safeer K. and Anbarasi B. (2010), Analytical method development and validation of Amlodipine and Hydrochlorothiazide in combined dosage form by RP-HPLC. Int. J. Chem.Tech. Res., 2(01), 21-25.

Lindsay S. (1991), HPLC, John Wiley and Sons, 30-45

Malaviya K. and Jat R. (2012), RP-HPLC Method Development and its Validation for Assay of Rupatadine Fumarate in Tablet Dosage Form, Inventi Rapid Pharm Ana & Qual Assurance, 3(4), 308.

McCarberg B. and Gibofsky A. (2012), Need to develop new nonsteroidal anti-inflammatory drug formulations, Clinical Therapeutics, 34 (9), 1954–63

Mendham J., Denny R. C. and Thomas M. (2004), In Vogel's Text Book of Quantitative Analysis; Pearson Education Limited, 6th Edn; 1-5.

Meyyanathan N., Muralidharan S., Rajan S., Gopal K., Suresh B. (2008), A simple sample preparation with HPLC-UV method for estimation of Amlodipine from plasma: Application to bioequivalence study, The Open Chemical and Biomedical Methods J. 2008, 1, 22-7.

Minson J. (2000), Pharmaceutical Analysis Dekker Series, 2(B), 16-77.

Oza K. (2011), Determination of losartan potassium and perindopril erbumine in combined tablet dosage form, Int. J. Pharm. and Bio. Sci., 2(01), 709-15.

Pachauri S., Paliwal S., Srinivas S. and Singh Y. (2010), Development & validation of HPLC method for analysis of some Antihypertensive agents in their pharmaceutical dosage forms, J. Phar. sc. and res., 2(08), 459-64.

Pandey R., Patil O., Bari B. and Dhumal M. (2014), Simultaneous estimation of Etodolac and Thiocolchicoside in bulk and in tablet formulation by UV-Spectrophotometry, Chem Ind Chem Eng Q, 20(1), 9-17.

Pattan R., Jamdar G., Godge K., Dighe S., Daithankar V. and Nirmal A. (2009), Journal of Chemical and Pharmaceutical Research, 1(1), 329-35.

Permender R., Sushila R., Shyama T. and Vikash K. (2010), Simultaneous estimation of Amlodipine besylate and Lisinopril dehydrate as A.P.I and in tablet dosage forms by modified form of simultaneous equation method using derivative uv spectroscopy, Int. J of PharmaTech research, 2(1), 556-62.

Prasada R., Rahaman A., Ragjendra Y. and Gangi P. (2010), RP-HPLC method for simultaneous estimation of Amlodipine besylate and Metaprolol in combined dosage form. IJPRD, 2(9), 69-76.

Rahman N. and Nasrul H. (2003), Validated spectrophotometric methods for the determination of Amlodipine besylate in drug formulations using

2,3dichloro5,6dicyano1,4 benzoquinone and ascorbic acid, J of pharmaceutical and biomedical analysis, 381-92.

Rahman N., Manisha S. and Nasrul H. (2004), Applications of oxidants to the spectrophotometric determination of Amlodipine besylate in pharmaceutical dosage forms. IL FARMACO, 59, 913-19.

Richards A., Harver P. and Champe C. (1995), Anti-inflammatory Drugs, unit VII ch.39, in Lipincott's Illustrated Reviews Pharmacology, 2nd edn. pp: 409.

Sankar D., Somasekhar V. and Shiva N. (2009), Development and validation of a rapid RP-HPLC method for the determination of venlafaxine hydrochloride in pharmaceutical dosage forms using experimental design. E-J. Chem., 6(04), 1091-102.

Sawant R., Bhangale L. and Joshi R. (2011), Journal of Chemical Metrology, 4(1), 21-7.

Shah A, Shah J., Shah H. and Upadhyay U. (2013), Formulation Development and Evaluation of Buccal Bilyer Patch using Thiocolchicoside and Diclofenac Sodium, Int J Inv Pharm Sci., 1(3), 190-211.

Shah V. (2001), Guidance for Industry, Bioanalytical Methods Validation.

Sharma K. (2002), In Instrumental methods of chemical analysis; Goel Publishing House, Krishna Prakashan Ltd, 19th Edn; 1- 4

Shukla A., Jat R., Sharma P. and Patel Y. (2011), Development and Validation of a Headspace Gas Chromatographic Method for Determination of Residual Solvents in Levetiracetam, Int J Pharm Sci Res, 2(5), 1270-75.

The United States Pharmacopeia XXII (1995), Validation of compendial Methods, USP Convention Inc. Rockville, MD, 95.

United States Pharmacopoeia (2014), Validation of Compendial Methods, Rockville MD USA, United States Pharmacopeial Convention Inc, USP 32 - NF 27, 1826-1827, 2103-2105.

Yuanyuan M., Feng Q., Xiaohong S., Xiumei L. and Famei L. (2007), Determination and pharmacokinetic study of Amlodipine in human plasma by ultra performance liquid chromatography-electrospray ionization mass spectroscopy. J of Pharm and Biomed Anal, 43, 1540-5.